



Dust Telescope on the Lunar Surface

E. Gruen, M. Horanyi, S. Auer, S. Robertson,
R. Srama, Z. Sternovsky


LASP, University of Colorado, Boulder, CO 80303
Physics Department, University of Colorado, Boulder, CO 80309
A&M Associates, Basye, VA, 22810

Overview

- Motivation: Science *from* the surface of the Moon
 - Detection of interstellar dust
 - Composition of interstellar dust
 - Interaction with the interplanetary magnetic field
- Instrumentation
- Summary

The Interstellar Dust Story (dust to dust)


Protoplanetary disk



Star + planetary system




Comets, asteroids




Interplanetary dust



Dust factories



Supernova
explosions




AGB stars



Diffuse interstellar medium




Stellar nebula



Formation of new stars




Dense molecular cloud



Ulysses: discovery of interstellar dust within the solar system


4



Interstellar dust identified by: Retrograde orbit, High impact velocity (> 26 km/s)

Properties of the interstellar dust flow

Flow direction




Frisch et al, 2007

ISD flux (at 1 AU)

$$\underline{J \approx 1.5 \times 10^{-4} \text{ m}^{-2} \text{ s}^{-1}}$$

(approx: 13 particles per m² per day)

Mass density distribution



The mass density distribution per log mass interval $mdn/d \ln(m)$ measured by Ulysses (Landgraf et al., 2000) shows an astonishing excess of large grains over the predictions of models based on wavelength-dependent extinction observations (continuous lines). This represents a challenge to our current understanding of the extinction measurements, as well as the chemistry of diffuse interstellar clouds.

Properties of the interstellar dust flow

Forces

$$F_L = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$$

\mathbf{B} - interplanetary

F_G - Sun's gravitational force ($\sim a^3$)

F_R - Radiation force ($\sim a^2$)

$$\beta = F_R / F_G$$


Cases

($r < 5$ nm) - tied to interstellar \mathbf{B}

($r < 200$ nm) - cannot reach Earth

($r \sim 300$ nm) - $\beta > 1$


Dust dynamics at 1 AU



$$m = 3 \times 10^{-13} \text{ g}$$

$$a = 0.3 \text{ } \mu\text{m}$$

$$\beta = 1.2$$




$$m = 1.2 \times 10^{-12} \text{ g}$$

$$a = 0.5 \text{ } \mu\text{m}$$

$$\beta = 0.8$$

Solar cycle effect

The change in the direction of ISD flow is due to solar cycle effects



Krueger et al, 2007

- Ulysses measures direction only by statistical means
- B field changes with solar cycle (22 years period)
- ~ 20 years flight time for dust to reach the Earth from the

Accurate trajectory measurement:



- Probes the IMF
- Provides source of ISD

ISD composition

- ISD grains contain most of the refractory elements (Fe, Mg, Si, Al, Ca)
- Significant content of C and O
- Organic material (PAH)
- Ices: H_2O , CO , CO_2 , CH_3OH (in dark clouds)
- **Mathis et al. (1996)**
 - a) Small graphite grains
 - b) Silicate grains
 - c) Composite grains containing carbon, silicates and oxides
- **Li & Greenberg (1997)**
 - a) Large grains with silicate core and organic refractory mantle
 - b) Small carbonaceous grains
 - c) Very small grains (or large clusters) of PAH
- **Draine & Li (2007)**
 - a) Amorphous silicate grains
 - b) Carbonaceous grains including small PAH

Stardust ISD sample return

Preliminary analysis started




- 195 days of sample collection
- Estimated 60 grains collected
- Heating and abrasion effects are likely severe
- Locating tracks and extraction are very challenging

Science objectives:

- **A1:** Explore the mass, chemical and isotopic composition distributions of ISD and Interplanetary (IDP) grains
- **A2:** Verify the Ulysses observations of the unexpectedly large flux of big ($m > 10^{-13}$ g) ISD grains
- **A3:** Validate remote sensing observations by in-situ measurements of the composition of ISD grains
- **A4:** Explore the connection between ancient ISD found in meteorites to that of 'fresh' inflowing ISD grains
- **H1:** Explore the influence of the solar cycle on the variability of the flow direction of ISD

In-situ Instrumentation: Dust Telescope



Dust Trajectory Sensor

Measures the velocity vector with high precision

Large Area Mass Analyzer



Measures the elemental composition

Target area: $\sim 0.1 \text{ m}^2$ (approx 1-2 impacts / day)

Z. Sternovsky

Dust Telescope prototypes


Large Area Mass Analyzer (LAMA)



Sternovsky et al, 2007


LAMA mass spectrum, 30 km/s Ni dust on Ag surface ($\underline{m/dm = 200}$)

Dust Trajectory Sensor (DTS)



Trajectory reconstruction, velocity < 1%, direction < 1° precision

Why from the Moon?



- Moon is a good platform for Dust Astronomy
- The number of ISD impacts on the Moon's surface is comparable to the impacts of interplanetary dust
- Instrumentations compatible with other dust detectors:
 - Interplanetary dust impacts
 - Secondary ejecta detectors
 - Slow moving dust

Summary

"All of us are truly and literally a little bit of stardust." – William A. Fowler

- Interstellar dust (ISD) impacts the Moon!
- Astrophysics and Heliophysics science goals
- Advanced instrumentation is ready



"SURE IT'S BEAUTIFUL, BUT I CAN'T HELP THINKING ABOUT THE INTERSTELLAR DUST OUT THERE."